



University of
Zurich^{UZH}

Zurich Open Repository and
Archive

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2020

Warm-up effect on handgrip strength in sedentary and overweight women

Hernández-Martínez, Jordan ; Rauch-Gajardo, Maria ; Cisterna, Diego ; Ramírez-Campillo, Rodrigo ; Moran, Jason ; Knechtle, Beat ; Nikolaidis, Pantelis Theodoros ; Álvarez, Cristian

Abstract: Introduction: The evaluation of handgrip strength has gained special relevance in the area of health. However, a standardized protocol of application is required to measure it, including warm-up procedures. Objective: To compare the acute effects of different warm-up strategies on maximal handgrip strength (MHS) in sedentary and overweight women. Materials and methods: Single-blind, randomized, cross-over study in which MHS was measured in 12 overweight women under the following conditions: i) no warm-up (control condition), ii) static stretching warm-up, iii) strength-based warm-up (i.e., resistance band exercise), and iv) isometric squeezing-ball warm-up for the forearm muscles. A Jamar dynamometer was used for the measurements, which were taken on four different days, at 48-hour rest intervals; three measurements were made per hand. Results: MHS mean values were 23.8 and 24.9 kg without warm-up, 20.3 and 21.4 kg after stretching warm-up, 20.9 and 22.9 kg after strength-based warm-up, and 22.0 and 23.0 kg after squeezing-ball warm-up for non-dominant and dominant hand, respectively. No significant ($p < 0.05$; one-way ANOVA) differences were observed between protocols, nor differences in MHS in relation to nutritional status, lean mass or fat mass. Conclusion: Warm-up is not required to measure MHS in overweight sedentary women when three measurements are made.

DOI: <https://doi.org/10.15446/revfacmed.v68n3.76057>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-190852>

Journal Article

Accepted Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Hernández-Martínez, Jordan; Rauch-Gajardo, Maria; Cisterna, Diego; Ramírez-Campillo, Rodrigo; Moran, Jason; Knechtle, Beat; Nikolaidis, Pantelis Theodoros; Álvarez, Cristian (2020). Warm-up effect on handgrip strength in sedentary and overweight women. *Revista Facultad de Medicina de la Universidad Nacional de Colombia*, 68(3):101-106.

DOI: <https://doi.org/10.15446/revfacmed.v68n3.76057>



Revista *de la* Facultad *de* Medicina

Publicación anticipada

Este artículo fue aprobado para publicación en el v68n3 de la Revista de la Facultad de Medicina teniendo en cuenta los conceptos de los pares evaluadores y los cambios realizados por los autores según estos conceptos. Por lo tanto, se publica la versión preliminar del artículo para su consulta y citación provisional, pero debe aclararse que esta puede diferir del documento final, ya que no ha completado las etapas finales del proceso editorial (corrección de estilo, traducción y diagramación) y solo los títulos, datos de autores, palabras clave y resúmenes corresponden a la versión final del artículo.

Esta versión puede consultarse, descargarse y citarse según se indique a continuación, pero debe recordarse que el documento final (PDF, HTML y XML) puede ser diferente.

Cómo citar:

Hernández-Martínez J, Rauch-Gajardo M, Cisterna D, Ramirez-Campillo R, Moran J, Knechtle B, et al. [Efecto del calentamiento en la fuerza de agarre de mano en mujeres sedentarias con sobrepeso]. Rev. Fac. Med. 2020;68(3): In press - 2020. English. doi: <http://dx.doi.org/10.15446/revfacmed.v68n3.76057>.

Article in press

This article was accepted for publication in V68N3 of Revista de la Facultad de Medicina (Journal of the Faculty of Medicine), considering the concepts of the peer reviewers and the changes made by the authors based on said concepts. Therefore, the preliminary version of this article is published for consultation and provisional citation purposes. However, it should be noted that this version may differ from the final document since it has not completed the final stages of the editorial process (proof-editing, translation, and layout). Only the titles, authorship, keywords and abstracts will remain unchanged in the final version of the article.

This version can be consulted, downloaded, and cited as indicated below, but please bear in mind that the final document (PDF, HTML, and XML) may differ.

How to cite:

Hernández-Martínez J, Rauch-Gajardo M, Cisterna D, Ramirez-Campillo R, Moran J, Knechtle B, et al. Effect of warm-up on hand grip strength in sedentary and overweight women. Rev. Fac. Med. 2020;68(3): In press - 2020. English. doi: <http://dx.doi.org/10.15446/revfacmed.v68n3.76057>.

Type of article: Original research**DOI:** <http://dx.doi.org/10.15446/revfacmed.v68n3.76057>**Title: Effect of warm-up on hand grip strength in sedentary and overweight women****Short title: Warm-up, handgrip strength and overweight****Título completo: Efecto del calentamiento en la fuerza de agarre de mano en mujeres sedentarias con sobrepeso****Título corto: Calentamiento, fuerza de agarre y sobrepeso.**

Received: 09/11/2018 Accepted: 04/02/2019

Jordan Hernández-Martínez^{1,2}, María Rauch-Gajardo², Diego Cisterna¹, Rodrigo Ramirez-Campillo¹, Jason Moran³, Beat Knechtle⁴, Pantelis Theodoros Nikolaidis⁵, Cristian Álvarez¹

¹ Universidad de Los Lagos - Department of Physical Activity Sciences - Human Performance Laboratory - Quality of Life and Wellness Research Group - Osorno - Chile.

² Universidad Santo Tomás - Occupational Therapy School - Faculty of Health - Osorno - Chile.

³ University of Essex - School of Sport, Rehabilitation and Exercise Sciences - Colchester - United Kingdom.

⁴ University of Zurich - Institute of Primary Care - Zurich - Switzerland.

⁵ University of West Attica - School of Health and Caring Sciences - Athens - Greece.

Corresponding author: Jordan Hernández-Martínez. Escuela de Terapia Ocupa-

cional, Facultad de Salud, Universidad Santo Tomás. Los Carrera No. 753, Edificio Central, Oficina General de la Escuela de Terapia Ocupacional. Phone: +56 9 95074360. Osorno. Chile. E-mail: jordan.eduardo.hernandez@gmail.com.

Abstract

Introduction: The evaluation of handgrip strength has gain special relevance in the area of health. However, a standardized protocol of application is required to measure it, including warm-up procedures.

Objective: To compare the acute effects of different warm-up strategies on maximal handgrip strength (MHS) in sedentary and overweight women.

Materials and methods: Single-blind, randomized, cross-over study in which MHS was measured in 12 overweight women under the following conditions: i) no warm-up (control condition), ii) static stretching warm-up, iii) strength-based warm-up (i.e., resistance band exercise), and iv) isometric squeezing-ball warm-up for the forearm muscles. A Jamar dynamometer was used for the measurements, which were taken on four different days, at 48-hour rest intervals; three measurement were made per hand.

Results: MHS mean values were 23.8 and 24.9 kg without warm-up, 20.3 and 21.4 kg after stretching warm-up, 20.9 and 22.9 kg after strength-based warm-up, and 22.0 and 23.0 kg after squeezing-ball warm-up for non-dominant and dominant hand, respectively. No significant ($p>0.05$; one-way ANOVA) differences were observed between protocols, nor differences in MHS in relation to nutritional status, lean mass or fat mass.

Conclusion: Warm-up is not required to measure MHS in overweight sedentary women when three measurements are made.

Keywords: Muscles; Body Fat; Women; Sarcopenia; Muscle Strength (MeSH).

Hernández-Martínez J, Rauch-Gajardo M, Cisterna D, Ramirez-Campillo R, Moran J, Knechtle B, et al. Effect of warm-up on hand grip strength in sedentary and overweight women. Rev. Fac. Med. 2020;68(3): - . English. doi: <http://dx.doi.org/10.15446/revfacmed.v68n3.76057>.

Resumen

Introducción. En los últimos años se ha dado una mayor importancia a la medición de la fuerza máxima de agarre de mano, sin embargo para hacer esta medición se requiere un protocolo estandarizado de aplicación, incluyendo procedimientos de calentamiento.

Objetivo. Comparar los efectos agudos de cuatro tipos de calentamiento en la fuerza máxima de agarre de mano de mujeres sedentarias con sobrepeso.

Materiales y métodos. Estudio ciego, aleatorizado y cruzado en el que se midió la fuerza máxima de agarre de mano de 12 mujeres con sobrepeso bajo las siguientes condiciones: i) sin calentamiento (condición de control), ii) con calentamiento de estiramiento estático, iii) con calentamiento basado en la fuerza (p. ej., ejercicios con banda elástica) y iv) con calentamiento con bola terapéutica de compresión para los músculos del antebrazo. Para las mediciones se utilizó un dinamómetro Jamar y estas se realizaron en cuatro días diferentes y en intervalos de 48 horas de descanso; además, se hicieron tres intentos de medición por mano.

Resultados. Los valores promedio de fuerza máxima de agarre para la mano no dominante y dominante fueron 23.8kg y 24.9kg sin calentamiento, 20.3kg y 21.4kg con estiramiento, 20.9kg y 22.9kg con banda elástica y 22.0kg y 23.0kg con bola terapéutica, respectivamente. No hubo diferencias significativas ($p > 0.05$; ANOVA de una vía) entre los protocolos, ni diferencias en la fuerza máxima de agarre de mano en relación con estado nutricional, masa magra o masa grasa.

Conclusión. No se requiere una sesión de calentamiento para medir la fuerza máxima de agarre de mano en mujeres sedentarias con sobrepeso cuando se realizan tres intentos de medición.

Palabras clave: Músculos; Mujer; Sarcopenia; Fuerza muscular (DeCS).

Hernández-Martínez J, Rauch-Gajardo M, Cisterna D, Ramirez-Campillo R, Moran J, Knechtle B, et al. [Efecto del calentamiento en la fuerza de agarre de mano en mujeres sedentarias con sobrepeso]. Rev. Fac. Med. 2020;68(3): - . English. doi: <http://dx.doi.org/10.15446/revfacmed.v68n3.76057>.

Introduction

Loss of muscle strength (dynapenia) (1) has a negative effect on morbidity and mortality (2). Therefore, the timely assessment of muscle strength is fundamental in preventive medicine (3).

The handgrip strength test is a validated and simple test used to assess muscle strength in several health-related contexts (4-10). Despite its importance in clinical practice, there is a wide range of choices regarding the equipment and protocols to use when measuring maximal handgrip strength (MHS) (11). Particularly, the effects of warming-up before performing a MHS tests have not yet been described.

A warm-up is generally intended to generate an increase in muscle temperature facilitating increased blood flow, optimizing metabolic responses (12-13), reducing the muscle's viscous resistance (i.e. smoother contraction) and increasing the speed of nerve transmission (14). By extension, the search for an optimal muscle temperature range that limits fatigue as much as possible whilst maximizing performance (12-15) seems prudent. Commonly, warm-up protocols tend to reflect the experience of individual researchers and practitioners, with most studies performed in athletes (16). Controlled studies regarding the effects of warm-up on maximal performance are particularly scarce, maybe due to the un-

willingness of voluntary subjects to complete a maximal effort without warm-up (i.e. control condition). Among the studies investigating the effect of warm-up protocols on muscular performance (e.g., maximal strength), conflicting results have arisen, with some studies showing an increase in performance after specific (17) or combined general and specific warm-up (18), while others have not (16).

Considering the lack of studies addressing the effects of warm-up on sedentary females with overweight and MHS, and the clinically relevant applicability of MHS in community-health programs (19), a standardized protocol of application is required. The aim of this study was to compare the acute effects of different warm-up strategies on MHS in sedentary females. It was hypothesized that different warm-up protocols would induce an effect on MHS.

Methods

Ethical approval

This study (study protocol N° 103- 2018) was approved by the Institutional Review Board of the Department of Physical Activity Sciences, Universidad de Los Lagos, as it is stated Minutes DECAF2016/3, issued on April 25, 2016. The participants who agreed to participate signed an informed consent form, after receiving an explanation of the risks and benefit derived from their participation in the study. The study was conducted according to the ethical principles of the latest version of the Declaration of Helsinki 2013 (20).

Subjects and procedures

By publicity in a local University, sedentary and overweight females ($n = 12$, age, 21.1 ± 2.0 y; $38.1\% \pm 8.4\%$ fat mass; further descriptive characteristics in table 1) were recruited in a single blind, randomized and cross-over study design. Participants completed four different measurement protocols to assess MHS, with 48 h of rest between each.

Table 1. Baseline characteristics of the sample.

Variables	Mean	SD
Body mass (kg)	64.5	9.1
Height (m)	158.3	8.4
Body mass index (kg/m ²)	26.3	3.9
Body fat (kg)	24.8	7.3
Lean mass (kg)	22.1	3.3
Water (L)	29.8	4.0
Lean mass left hand (kg)	2.1	0.4
Lean mass right hand (kg)	2.1	0.4
Fat mass left hand (kg)	1.8	0.7
Fat mass left hand (kg)	1.8	0.7

Abbreviations: kg: kilograms; m: meter; kg/m²: kilograms by square meter; L: liters.

Source: Own elaboration based on the data obtained in the study.

To be included in the study, participants were required to: i) be over 18 years old, ii) be sedentary (weekly physical activity level = 600 METs.min.week) (21), iii) be free of cardiovascular, pulmonary or skeletal muscle pathologies (22), iv) have a fat mass >30% of total body mass. All experimental procedures were performed under controlled and standardized conditions in the Laboratory of Human Performance at the University where the study was conducted, always at the same time of day, temperature, humidity, rest time (i.e., sleep hours before testing), menstrual cycle phase, and hours post last meal. Subjects' height (Bodimeter 206, SECA, Germany to 0.1 cm), body mass and body composition (InBody120, tetrapolar 8-point tactile electrodes system, model BPM040S12F07, Biospace, Inc., USA, to 0.1 kg) were measured according to previous recommendations (23).

Measurement of handgrip strength

The test was applied according to previous recommendations (24). To assess MHS, an adjustable digital dynamometer was used (Jamar®, PLUS+, Sammons

Preston, Patterson Medical, Illinois, United States). After randomly assigning the order of dominant and non-dominant hand assessment, subjects had three valid attempts to achieve MVIHS for both dominant and non-dominant hands, with 2 minutes of rest between attempts. For each maximal attempt, subjects were asked to exert 5 seconds of maximal effort, while receiving standardized verbal motivation. Subjects completed each maximal attempt while seated on a chair in an erect position. The hip, knee, and elbow were flexed to a 90° angle and the shoulder was abducted and neutrally rotated. The forearm was in a neutral position and the wrist was slightly extended (0° to 30°). Subjects performed the test with a horizontal cylinder handgrip using the position II on the digital dynamometer, while the evaluator lightly supported this in its base. The best result (expressed in kg) of the three valid maximal attempts for each hand was chosen for statistical analysis.

Warm-up protocols

Four randomly selected warm-up protocols (Table 2) were applied for the forearm muscles of both the dominant and non-dominant hands as follow: i) no warm-up (control condition), during which subjects remained seated comfortably for three minutes before testing; ii) static stretching warm-up which subjects carried out static stretching of the forearms flexors and extensors muscles for a total of five sets of five seconds each (25); iii) strength (i.e. elastic band-based) warm-up during which subjects completed two sets of ten repetitions for the forearm flexor muscles for a duration of 2.5 seconds for each contraction (26) using an elastic band (THERA Band™; medium intensity, blue color) and 30 seconds to one minute of recovery between sets; and iv) isometric therapeutic squeezing-ball warm-up during which subjects completed one grip per 2.5 seconds (for a total of twenty repetitions) on a therapeutic squeeze ball (27). The ten point Borg scale of subjective perception of effort was used to measure the intensity during warm-ups to standardize intensity across all conditions, always between 3 and 6

points. After the warm-ups, three minutes elapsed before MHS testing.

Table 2. Characteristics of the warm up protocols.

Warm up	Exercises	Sets	Repetitions	Recovery between sets	Recovery after warm up
No warm-up	-	-	-	-	-
Static stretching	Static flexion of wrist	5	5 seconds	30 seconds	3 minutes
	Static extension of wrist	5	5 seconds	30 seconds	
Elastic band	Dynamic flexion of wrist	2	10	30 seconds	3 minutes
Isometric therapeutic squeezing-ball	Squeeze and release	1	20	30 seconds	3 minutes

Note: the table is an elaboration of the authors, based on the data of the present study.

Statistical analysis

All values are reported as means \pm standard deviations. The Shapiro-Wilk and Levene's tests yielded non-significant values for all data. To determine the effects of the conditions on MHS absolute mean differences between conditions were compared using a repeated measures analysis of variance, with Fisher post hoc procedures. The α level was set at $p < 0.05$ for statistical significance, with Cohen's d representing effect size (ES) interpreted as <0.2 = trivial; $0.2-0.6$ = small; $>0.6-1.2$ = moderate; $>1.2-2.0$ = large; $>2.0-4.0$ = very large; >4.0 =

extremely large). The reliability of the assessments was determined using the intra-class correlation coefficient. All measurements yield values ≥ 0.9 .

Results

The MHS mean values for the non-dominant and dominant hand were, 23.8 kg and 24.9 kg after no warm-up, 20.3 kg and 21.4 kg after the stretching warm-up, 20.9 kg and 22.9 kg after the strength warm-up, and 22.0 kg and 23.0 kg after the squeezing-ball warm-up, respectively (Figure 1). No significant differences ($p > 0.05$; $ES < 0.2$) were observed among warm-up protocols.




Figure 1. Abbreviations: Lh: left hand; Rh: right hand; Sav: Denotes mean maximal strength values from three measurement attempts; Smax: Denotes maximal strength value from three measurement attempts; Cal-EB: strength warm-up; Cal-Flex: stretching warm-up; Cal-Ball: squeezing-ball warm-up; w/ Cal: no warm-up; kg: kilograms. NS: Denotes non-significant differences within-groups and between groups. Note: the figure is an elaboration of the authors, based on the data of the present study.

For the maximal strength value obtained from three MHS attempts (S_{max} in figure 1) and the mean strength value obtained from three MHS attempts (S_{av} in figure 1), no significant differences were observed between values ($p > 0.05$; $ES < 0.2$).

Discussion

The aim of this study was to compare the effects of different warm-up protocols on MHS. Main findings suggest that the three randomly selected warm-up protocols had no effect on MHS in a sample of 12 sedentary and overweight females. Moreover, a reduced MHS trend was observed in the participants after performing a static stretching based warm-up.

Regarding static stretching in warm-up routines, Behm et al. (28), in a study about the effects of static stretching warm-up on the strength of quadriceps muscles, reported a significant 12% maximal isometric strength decrease. Similar results have also been described for the pectoralis major and the triceps brachii muscles (29). In this sense, the results reported in the current study are in agreement with the aforementioned findings (28-29), since forearm flexor and extensor muscles static stretching, regardless of hand dominance, negatively affected MHS in sedentary and overweight females. Several factors may help to explain the impairment in MHS after static stretching, such as alterations in the mechanical components of muscle contraction (30), decreased muscle activation (28), or both (30).

In the current study, compared to the control condition, there were no improvements in MHS after the warm up with elastic band. This finding is in contrast with the results of a study conducted by Tilley and Macfarlane (31), where an increase in swing performance was demonstrated in elite male golfers after a warm-up with a rubber band. In male judokas, a warm-up with an elastic band allow them to improve the high jerk test performance when compared to a control condition (32). Moreover, Mina et al. (33), observed an increase in maximal

squat strength in men after a warm up with an elastic band. In addition, in a study conducted by Aandahl et al. (34) an increase in the maximal kick speed in martial arts fighters was observed leading the authors to conclude that this increase was due to greater recruitment of higher order motor units, greater synchronization of the motor units and low presynaptic inhibition. However, the performance-enhancing factors observed in previous studies (31-34) were found in athletes, not in a sedentary population as in the present study. Notably, the aforementioned studies (31-34) usually analyzed the effect of elastic band warm ups on large muscle groups in multi-joint exercise, contrasting to the muscle groups analyzed in our study. Therefore, these methodological elements (i.e., sedentary vs. athletes; small muscle group vs. large muscle group; single-joint vs. multi-joint) could help explain the different results found in this work and those previously published (31-34).

Current results shows that the specific warm up with a therapeutic ball (squeezing-ball warm-up) had no effect on MHS when compared to the control condition. . A specific warm up involves skill exercises which demonstrate equivalency with the targeted motor task (35). It seeks to increase performance (36) via increases in muscle temperature, reductions in muscle viscosity resistance and greater nerve transmission speed (37). In a study conducted by Andrade et al. (38), the effects of a general warm up, a specific warm up and a combined warm up on explosive muscular performance were compared, with improvements in squat jump and drop jump being observed after a specific jump-based warm up. Similarly, in a study conducted in volleyball players, an improvement in counter-movement vertical jump was observed after a specific warm up protocol based on jump exercises (39). Of note, the improvements in jumping performance after specific jump-based warm ups were observed in large muscle groups. Smaller muscles, such as the forearm, are composed of a significant number of slow-twitch muscle fibers that require a low motor unit firing frequency (i.e., 5 to 30 Hz), unlike other larger muscle groups (40). Such slow-twitch fibers are easily

excitable (41) and so require lower levels of stimulation to achieve maximal activation and, therefore, maximal strength. Therefore, as muscular forearm activation in hand-grip tasks is relatively easier (42) as compared to larger muscle groups, a specific warm-up may not add to the performance of such muscle group during hand-grip tasks.

Of note, no differences were observed in MHS after dynamic (i.e., elastic band) and isometric (i.e., static stretching; isometric therapeutic squeezing-ball) warm-up protocols. Such observation seems to be in contrast with the findings of a previous study (43), where a dynamic warm up, when compared to static-stretching warm up, improved power and agility (T-shuttle run, underhand medicine ball throw for distance, and 5-step jump) in male and females military cadets. However, it should be acknowledge the methodological differences between studies, such as the participant's characteristics (i.e., females vs. mix sample of male and females), physical fitness level (i.e., low vs. high), type of performance test (i.e., maximal isometric strength vs. dynamic power test), among others. In this regard, the American College of Sport Medicine indicated that more controlled studies are needed to substantiate the effectiveness of warm-up protocols (44). The lack of consensus may be partially related to the different methodological issues previously indicated, as the effect of warm-up may vary according to such elements (45). Moreover, most studies related to warm-up strategies have been conducted in athletes (45). In this sense, the present results expand the limited knowledge available about the effect of different warm up protocols on the MHS in sedentary and overweight females.

Limitations, strength and practical applications

A limitation of the study was the sample size, as it may not have allowed statistically significant findings to be observed. Future studies may aim to replicate the current findings with a greater sample size. Additionally, in order to better understand the underlying mechanisms of different warm-up protocols, future

research may include biomechanical as well as physiological measures related to the responses of forearm muscles to different warm-up protocols in sedentary and overweight females.

Conclusion

Warm-up of the forearm muscles does not acutely increase isometric MHS in sedentary overweight females in the dominant, or the non-dominant hands. Three isometric attempts, without warm-up, allows the achievement of MHS, with high reliability, serving as a time-efficient measurement protocol, with high applicability in clinical assessment.

Conflict of interest

This study did not have any conflict of interest.

Financing

None declared by the authors.

References

- 1 Clark BC & Manini TM. What is dynapenia? *Journal Nutrition*. 2012; 28 (5). 495-503.
- 2 Gale CR, Martyn CN, Cooper C, Sayer AA. Grip strength, body composition, and mortality. *International journal of epidemiology* 2007; 36(1): 228-35.
- 3 Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, Martin FC, Michel JP, Rolland Y, Schneider SM, Topinková E, Vandewoude M, Zamboni M; European Working Group on Sarcopenia in Older People. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age Ageing*. 2010;39(4):412-23.
- 4 Abizanda P, Navarro JL, García-Tomás MI, López-Jiménez E, Martínez-Sán-

chez E, Paterna G. Validity and usefulness of hand-held dynamometry for measuring muscle strength in community-dwelling older persons. *Archives of Gerontology and Geriatrics*. 2012; 54 (1). 21-27.

5 Cuesta A, Hilgenkamp T. Reference Values of Grip Strength Measured with a Jamar Dynamometer in 1526 Adults with Intellectual Disabilities and Compared to Adults without Intellectual Disability. *PLoS ONE*. 2015; 10 (6). e0129585.

6 Kim S, Lee H, Cho E. Low Handgrip Strength is Associated with Low Bone Mineral Density and Fragility Fractures in Postmenopausal Healthy Korean Women. *Journal of Korean Medical Science*. 2012; 27 (7). 744-747.

7 Botoseneanu A, Bennett J, Nyquist L, Shinkai S, Fujiwara Y, Yoshida H, Aiello A, Cigolle, Ch, Liang J. Cardiometabolic Risk, Socio-Psychological Factors, and Trajectory of Grip Strength Among Older Japanese Adults. *Journal of Aging and Health*. 2015; 27 (7). 1123-46.

8 Leong D, Teo K, Rangarajan S, Lopez P, Avezum A, Orlandini A, Seron P, Ahmed S, Rosengren A, Kelishadi R, Rahman O, Swaminathan S, Iqbal R, Gupta R, Lear S, Oguz A, Yusoff K, Zatonska K, Chifamba J, Igumbor E, Mohan V, Anjana R, Gu H, Li W, Yusuf S. Prognostic value of grip strength: findings from the Prospective Urban Rural Epidemiology (PURE) study. *Lancet*; 2015; 386 (9990). 266-273.

9 Clifford M, Hamer P, Phillips M, Wood F, Edgar D. Grip strength dynamometry: Reliability and validity for adults with upper limb burns. *Burns*. 2013; 39 (7). 1430-1436.

10 Di Monaco M, Castiglioni C, De Toma E, Gardin L, Giordano S, Tappero R. Handgrip strength is an independent predictor of functional outcome in hip-fracture women: a prospective study with 6-month follow-up. *Medicine (Baltimore)*. 2015; 94(6): e542.

- 11 Roberts H, Denison H, Martin H, Patel H, Syddall H, Cooper C, Sayer A. A review of the measurement of grip strength in clinical and epidemiological studies: towards a standardised approach. *Age and Ageing* 2011; 40 (4): 423-429.
- 12 Bishop D. Warm up I: potential mechanisms and the effects of passive warm up on exercise performance. *Sports Med.* 2003; 33(6):439-54.
- 13 McGowan CJ, Pyne DB, Thompson KG, et al. Warm-up strategies for sport and exercise: mechanisms and applications. *Sports Med.* 2015; 45:1523-46.
- 14 Woods K, Bishop P, Jones E. Warm-up and stretching in the prevention of muscular injury. *Sports Med* 2007; 37(12): 1089-99.
- 15 Racinais S, Oksa J. Temperature and neuromuscular function. *Scand J Med Sci Sports.* 2010; 20(3):1-18.
- 16 Blazeovich AJ, Gill ND, Kvorning T, Kay AD, Goh AG, Hilton B, Drinkwater EJ, Behm DG. No Effect of Muscle Stretching within a Full, Dynamic Warm-up on Athletic Performance. *Med Sci Sports Exerc.* 2018; 50(6):1258-1266.
- 17 Burkett LN, Phillips WT, Ziuraitis J. The best warm-up for the vertical jump in college-age athletic men. *J Strength Cond Res* 2005; 19(3): 673-6.
- 18 Abad CC, Prado ML, Ugrinowitsch C, Tricoli V, Barroso R. Combination of general and specific warm-ups improves leg-press one repetition maximum compared with specific warm-up in trained individuals. *J Strength Cond Res* 2011; 25(8): 2242-5.
- 19 Garcia-Hermoso A, Cofre-Bolados C, Andrade-Schnettler R, Ceballos-Ceballos R, Fernández-Vergara O, Vegas-Heredia ED, Ramírez-Vélez R, Izquierdo M. Normative Reference Values for Handgrip Strength in Chilean Children at 8-12 Years Old Using the Empirical Distribution and the Lambda, Mu, and Sigma Statistical Methods. *J Strength Cond Res.* 2018; doi: 10.1519/JSC.0000000000002631.

- 20 Declaracion de Helsinki de la Asociacion Medica Mundial. Principios éticos para las investigaciones médicas en seres humanos, 64ª Asamblea General, Fortaleza, Brasil, octubre 2013; (accessed www.wma.net).
- 21 IPAQ. Research Committe. Guidelines for the data processing and analysis of the International Physical Activity Questionnaire 2005; Available from: www.ipaq.ki.se.
- 22 PAR-Q. The Physical Activity Readiness Questionnaire: PAR-Q & YOU. 2002 (accessed www.csep.ca/forms).
- 23 Marfell M, Olds T, Stewart A, Carte L. International Standards for Anthropometric Assessment. Potchefstroom, South Africa. 2006.
- 24 Fess, E, Moran Ch. American Society of Hand Therapists Clinical Assessment Recommendations. 1981.
- 25 Waymel T, Choque, J. 250 Ejercicios de estiramiento y tonificación muscular. 3º Edicion. España: Editorial Paidotrivo; 2009.
- 26 Lopez M, Lozano V, Yuguero A, Fontseré N. Influencia del ejercicio físico en el desarrollo de fistulas arteriovenosas nativas. Enfermeria Neurologica 2015; 18 (3). 168-73.
- 27 Luginbühl R, Brunner F, Schneeberger AG. No effect of forearm band and extensor strengthening exercises for the treatment of tennis elbow: a prospective randomized study. Chir Organi Mov 2008; 91(1): 35-40.
- 28 Behm D, Button D, & Butt J. Factors affecting force loss with prolonged stretching. Canadian Journal of Applied Physiology, 2001; 26: 262-72.
- 29 Leone DC, Pezarat P, Valamatos MJ, Fernandes O, Freitas S, Moraes AC. Upper body force production after a low-volume static and dynamic stretching. Eur J Sport Sci 2014; 14 (1). 69-75.

- 30 Cramer JT, Beck TW, Housh TJ, Massey LL, Marek SM, Danglemeier S, Purkayastha S, Culbertson JY, Fitz KA, Egan AD. Acute effects of static stretching on characteristics of the isokinetic angle - torque relationship, surface electromyography, and mechanomyography. *J Sports Sci.* 2007; 25 (6): 687-98.
- 31 Tilley NR, Macfarlane A. Effects of different warm-up programs on golf performance in elite male golfers. *Int J Sports Phys Ther.* 2012;7(4):388-95.
- 32 Lum D. Effects of Various Warm Up Protocol on Special Judo Fitness Test Performance. *J Strength Cond Res.* 2017. DOI: 10.1519/JSC.0000000000001862.
- 33 Mina MA, Blazeovich AJ, Giakas G, Kay AD. Influence of variable resistance loading on subsequent free weight maximal back squat performance. *J Strength Cond Res.* 2014;28(10):2988-95.
- 34 Aandahl HS, Van den Tillaar, R, and Von Heimburg, E. Effect of post-activation potentiation on kinematics and kicking performance in a roundhouse kick with trained martial arts practitioners. *Proceedings of the 33rd International Conference on Biomechanics in Sports.* Colloud, F, Domalain, M, Monnet, T, eds. Poitiers, France, 2015. pp. 281-284.
- 35 Young WB, Behm D.G. Should Static Stretching Be Used During a Warm-Up for Strength and Power Activities? *Strength Cond J.* 2002;24(6):33-37.
- 36 Thompsen AG, Kackley T, Palumbo MA, Faigenbaum AD. Acute effects of different warm-up protocols with and without a weighted vest on jumping performance in athletic women. *J Strength Cond Res.* 2007;21(1):52-56.
- 37 Woods K, Bishop P, Jones E. Warm-up and stretching in the prevention of

muscular injury. Sports Med. 2007;37(12):1089-1099.

38 DC Andrade, C Henriquez-Olguín, AR Beltrán, MA Ramírez, C Labarca, M Cornejo, C Álvarez, R Ramírez-Campillo. Effects of general, specific and combined warm-up on explosive muscular performance. Biol Sport. 2015; 32(2): 123-128.

39 Rezende Fernando Nazário de, Mota Gustavo Ribeiro da, Lopes Charles Ricardo, Silva Bruno Victor C. da, Simim Mário A. M., Marocolo Moacir. Specific warm-up exercise is the best for vertical countermovement jump in young volleyball players. Motriz: rev. educ. fis. 2016; 22(4): 299-303.

40 Astrand P, Rodahl K, Dahl H, Stromme S. Funcion motora. González P. Manual de fisiología del ejercicio. 4º Edición. Badalona (España): Editorial Paidotribo; 2010. 89-153.

41 Henneman E, Somjen G, Carpenter DO. Excitability and inhibibility of motoneurons of different sizes. Journal of Neurophysiology, 1965; 28: 599-620.

42 Fattorini L, Tirabasso A, Lunghi A, Di Giovanni R, Sacco F, Marchetti E. Muscular forearm activation in hand-grip tasks with superimposition of mechanical vibrations. J Electromyogr Kinesiol. 2016; 26:143-8.

43 McMillian DJ, Moore JH, Hatler BS, Taylor DC. Dynamic vs. static-stretching warm up: the effect on power and agility performance. J Strength Cond Res. 2006; 20:492-9.

44 Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, et al. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. Med Sci Sports Exerc. 2011; 43:1334-59.

45 McGowan CJ, Pyne DB, Thompson KG, Rattray B. Warm-up strategies for sport and exercise: Mechanisms and applications. Sports Med. 2015; 45:1523-46.

Publicación anticipada.
Rev. Fac. Med.